

# ConvolutEd Mathematics: A Comprehensive Study

Pu Justin Scarfy Yang

August 03, 2024



# Contents

<b>1</b>	<b>Introduction</b>	<b>5</b>
1.1	Overview . . . . .	5
1.2	Objectives . . . . .	5
<b>2</b>	<b>Convolved Linear Algebra</b>	<b>7</b>
2.1	Convolved Matrix Theory . . . . .	7
2.1.1	Definitions . . . . .	7
2.1.2	Properties . . . . .	7
2.1.3	Example: Convolved 2x2 Matrix . . . . .	7
2.2	Convolved Eigenvalues and Eigenvectors . . . . .	8
2.2.1	Definitions . . . . .	8
2.2.2	Theorem on Eigenvalue Properties . . . . .	8
2.2.3	Example: Convolved Eigenvalues of a Matrix . . . . .	8
<b>3</b>	<b>Convolved Differential Equations</b>	<b>9</b>
3.1	Convolved Ordinary Differential Equations (ODEs) . . . . .	9
3.1.1	Definitions . . . . .	9
3.1.2	Theorem on Existence and Uniqueness . . . . .	9
3.1.3	Example: Solving a Convolved ODE . . . . .	9
3.2	Convolved Partial Differential Equations (PDEs) . . . . .	10
3.2.1	Definitions . . . . .	10
3.2.2	Theorem on Solvability . . . . .	10
3.2.3	Example: Heat Equation . . . . .	10
<b>4</b>	<b>Convolved Topological Structures</b>	<b>11</b>
4.1	Convolved Metric and Topological Spaces . . . . .	11
4.1.1	Definitions . . . . .	11
4.1.2	Theorem on Open Sets . . . . .	11

4.1.3	Example: Convoluted Open Sets in $\mathbb{R}$ . . . . .	12
4.2	Convoluted Manifolds . . . . .	12
4.2.1	Definitions . . . . .	12
4.2.2	Theorem on Transition Maps . . . . .	12
4.2.3	Example: Convoluted Manifolds in $\mathbb{R}^2$ . . . . .	12
<b>5</b>	<b>Advanced Convoluted Homotopy and Cohomology</b>	<b>13</b>
5.1	Convoluted Homotopy Groups . . . . .	13
5.1.1	Definitions . . . . .	13
5.1.2	Theorem on Homotopy Equivalence . . . . .	13
5.1.3	Example: Convoluted Homotopy of $S^1$ . . . . .	13
5.2	Convoluted Cohomology Theories . . . . .	14
5.2.1	Definitions . . . . .	14
5.2.2	Theorem on Exact Sequences . . . . .	14
5.2.3	Example: Convoluted Cohomology of a Torus . . . . .	14
<b>6</b>	<b>Convoluted Logic and Set Theory</b>	<b>15</b>
6.1	Convoluted Logical Systems . . . . .	15
6.1.1	Definitions . . . . .	15
6.1.2	Theorem on Logical Consistency . . . . .	15
6.1.3	Example: Convoluted Logical Expressions . . . . .	16
6.2	Convoluted Set Theory . . . . .	16
6.2.1	Definitions . . . . .	16
6.2.2	Theorem on Set Operations . . . . .	16
6.2.3	Example: Convoluted Power Set . . . . .	16
<b>7</b>	<b>Future Directions and Research</b>	<b>17</b>
7.1	Potential Applications . . . . .	17
7.2	Ongoing Research . . . . .	17
<b>8</b>	<b>Appendices</b>	<b>19</b>
<b>A</b>	<b>Appendix A: Detailed Proofs</b>	<b>21</b>
<b>B</b>	<b>Appendix B: Supplementary Material</b>	<b>23</b>

# Chapter 1

## Introduction

### 1.1 Overview

This book introduces and rigorously develops the concept of convoluted mathematics. By modifying classical mathematical structures and operations using a convolution function  $\gamma$ , we explore new perspectives and applications across various mathematical domains.

### 1.2 Objectives

- Define and analyze convoluted structures in linear algebra, differential equations, topology, and logic. - Develop and prove theorems related to convoluted operations. - Provide examples and applications in diverse mathematical contexts.



# Chapter 2

## Convolved Linear Algebra

### 2.1 Convolved Matrix Theory

#### 2.1.1 Definitions

A convolved matrix  $\mathbf{A}_\gamma$  is defined by applying  $\gamma$  to the entries of a matrix  $\mathbf{A}$ . For a matrix  $\mathbf{A} = [a_{ij}]$ , the convolved matrix is:

$$\mathbf{A}_\gamma = [\gamma(a_{ij})]$$

#### 2.1.2 Properties

##### Determinant

If  $\mathbf{A}$  is invertible,  $\mathbf{A}_\gamma$  is invertible, and:

$$\det_\gamma(\mathbf{A}^{-1}) = \frac{1_\gamma}{\det_\gamma(\mathbf{A})}$$

Show  $\gamma$  preserves invertibility and multiplicative identity. Using  $\gamma$ 's homomorphic properties ensures that:

$$\det(\mathbf{A} \cdot \mathbf{A}^{-1}) = \det(\mathbf{I}) \Rightarrow \det_\gamma(\mathbf{A}_\gamma \cdot \mathbf{A}_\gamma^{-1}) = 1_\gamma$$

#### 2.1.3 Example: Convolved 2x2 Matrix

Consider  $\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . The convolved matrix is:

$$\mathbf{A}_\gamma = \begin{bmatrix} \gamma(a) & \gamma(b) \\ \gamma(c) & \gamma(d) \end{bmatrix}$$

Its convoluted determinant is:

$$\det_\gamma(\mathbf{A}) = \gamma(ad - bc)$$

## 2.2 Convoluted Eigenvalues and Eigenvectors

### 2.2.1 Definitions

For a square matrix  $\mathbf{A}_\gamma$ , the convoluted eigenvalue  $\lambda_\gamma$  and eigenvector  $\mathbf{v}_\gamma$  satisfy:

$$\mathbf{A}_\gamma \mathbf{v}_\gamma = \lambda_\gamma \mathbf{v}_\gamma$$

### 2.2.2 Theorem on Eigenvalue Properties

If  $\lambda$  is an eigenvalue of  $\mathbf{A}$ , then  $\lambda_\gamma = \gamma(\lambda)$  is an eigenvalue of  $\mathbf{A}_\gamma$ .

Apply  $\gamma$  to the eigenvalue equation  $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ . Using  $\gamma$ 's linearity:

$$\gamma(\mathbf{A}\mathbf{v}) = \gamma(\lambda\mathbf{v}) \Rightarrow \mathbf{A}_\gamma \mathbf{v}_\gamma = \lambda_\gamma \mathbf{v}_\gamma$$

### 2.2.3 Example: Convoluted Eigenvalues of a Matrix

Consider a matrix  $\mathbf{A}$  with eigenvalues  $\lambda_1, \lambda_2$ . The convoluted eigenvalues are  $\lambda_{\gamma,1} = \gamma(\lambda_1)$  and  $\lambda_{\gamma,2} = \gamma(\lambda_2)$ .



# Chapter 3

## Convolved Differential Equations

### 3.1 Convolved Ordinary Differential Equations (ODEs)

#### 3.1.1 Definitions

A convolved ODE is defined as:

$$\gamma\left(\frac{dy}{dt}\right) = \gamma(f(t, y))$$

#### 3.1.2 Theorem on Existence and Uniqueness

If  $f$  is Lipschitz continuous and  $\gamma$  preserves differentiability, the convolved ODE has a unique solution.

Use the Picard-Lindelöf theorem and show  $\gamma$  preserves contraction properties. Specifically, for any two solutions  $y_1, y_2$ :

$$\|\gamma(y_1(t)) - \gamma(y_2(t))\| \leq L\|\gamma(y_1) - \gamma(y_2)\|$$

#### 3.1.3 Example: Solving a Convolved ODE

Consider the ODE  $\frac{dy}{dt} = y$ . Its convolved form is:

$$\gamma\left(\frac{dy}{dt}\right) = \gamma(y)$$

Solution is given by  $\gamma(y(t)) = \gamma(Ce^t)$ , where  $C$  is a constant.

## 3.2 Convoluted Partial Differential Equations (PDEs)

### 3.2.1 Definitions

A convoluted PDE is defined as:

$$\gamma\left(\frac{\partial u}{\partial t}\right) = \gamma(\nabla^2 u)$$

### 3.2.2 Theorem on Solvability

If  $\gamma$  is linear and preserves boundary conditions, the convoluted PDE has solutions mirroring the original PDE.

Demonstrate existence through separation of variables and Fourier transforms. Ensure that boundary conditions are met by applying  $\gamma$  to both the PDE and the conditions:

$$\gamma(u(x, 0)) = \gamma(g(x))$$

### 3.2.3 Example: Heat Equation

Consider the heat equation  $\frac{\partial u}{\partial t} = \nabla^2 u$ . The convoluted form is:

$$\gamma\left(\frac{\partial u}{\partial t}\right) = \gamma(\nabla^2 u)$$

Solve using convoluted separation of variables to obtain:

$$u_\gamma(x, t) = \sum_{n=1}^{\infty} \gamma(a_n e^{-\gamma(\lambda_n)t} \phi_n(x))$$

# Chapter 4

## Convolutated Topological Structures

### 4.1 Convolutated Metric and Topological Spaces

#### 4.1.1 Definitions

A convolutated metric space  $(X_\gamma, d_\gamma)$  is defined by:

$$d_\gamma(x, y) = \gamma(d(x, y))$$

A convolutated topology  $\tau_\gamma$  is:

$$\tau_\gamma = \{\gamma(U) \mid U \in \tau\}$$

#### 4.1.2 Theorem on Open Sets

If  $(X, \tau)$  is a topological space and  $\gamma$  preserves set operations,  $(X_\gamma, \tau_\gamma)$  is topological.

Verify open set axioms under  $\gamma$ . For any open sets  $U, V \in \tau$ ,

$$\gamma(U \cup V) = \gamma(U) \cup \gamma(V)$$

and

$$\gamma(U \cap V) = \gamma(U) \cap \gamma(V)$$

show preservation of union and intersection.

### 4.1.3 Example: Convoluted Open Sets in $\mathbb{R}$

Consider open intervals in  $\mathbb{R}$ :  $(a, b)$ . The convoluted open interval is:

$$(a_\gamma, b_\gamma) = \{\gamma(x) \mid a < x < b\}$$

## 4.2 Convoluted Manifolds

### 4.2.1 Definitions

A convoluted manifold  $M_\gamma$  is a manifold  $M$  where the smooth structure is modified by  $\gamma$ . Charts are transformed as:

$$\varphi_\gamma : U \rightarrow \gamma(\mathbb{R}^n)$$

**Notation:** Use  $\mathcal{M}_\gamma$  for convoluted manifolds.

### 4.2.2 Theorem on Transition Maps

Transition maps  $\varphi_\gamma \circ \psi_\gamma^{-1}$  are smooth if  $\gamma$  preserves smoothness.

Show  $\gamma$  retains differentiability of transition functions. If  $\varphi \circ \psi^{-1}$  is smooth, then

$$\varphi_\gamma \circ \psi_\gamma^{-1} = \gamma(\varphi \circ \psi^{-1})$$

is smooth by  $\gamma$ 's property.

### 4.2.3 Example: Convoluted Manifolds in $\mathbb{R}^2$

Consider a circle  $S^1$ . Its convoluted chart is:

$$\varphi_\gamma(\theta) = (\gamma(\cos(\theta)), \gamma(\sin(\theta)))$$

Verify smooth transitions between charts via  $\gamma$ .

# Chapter 5

## Advanced Convolutated Homotopy and Cohomology

### 5.1 Convolutated Homotopy Groups

#### 5.1.1 Definitions

A convolutated homotopy group  $\pi_n(X)_\gamma$  is defined as:

$$\pi_n(X)_\gamma = \gamma(\pi_n(X))$$

**Notation:**  $\pi_\gamma$  for convolutated homotopy groups.

#### 5.1.2 Theorem on Homotopy Equivalence

If  $X \sim Y$ , then  $X_\gamma \sim Y_\gamma$ .

Preserve homotopy equivalence under  $\gamma$ . For a homotopy  $H : X \times I \rightarrow Y$ ,

$$\gamma(H) : \gamma(X) \times \gamma(I) \rightarrow \gamma(Y)$$

ensures  $\gamma$  maps homotopies to homotopies.

#### 5.1.3 Example: Convolutated Homotopy of $S^1$

For  $S^1$  with  $\pi_1(S^1) \cong \mathbb{Z}$ , the convolutated homotopy group is:

$$\pi_1(S^1)_\gamma \cong \gamma(\mathbb{Z})$$

## 5.2 Convoluted Cohomology Theories

### 5.2.1 Definitions

A convoluted cohomology theory  $H_\gamma^n(X)$  is given by:

$$H_\gamma^n(X) = \gamma(H^n(X))$$

**Notation:**  $H_\gamma$  for convoluted cohomology groups.

### 5.2.2 Theorem on Exact Sequences

Convoluted exact sequences remain exact if  $\gamma$  is exact-preserving.

Show sequence exactness remains under  $\gamma$ . For any exact sequence:

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

$\gamma$  ensures:

$$0_\gamma \rightarrow \gamma(A) \rightarrow \gamma(B) \rightarrow \gamma(C) \rightarrow 0_\gamma$$

### 5.2.3 Example: Convoluted Cohomology of a Torus

For a torus  $T^2$  with  $H^1(T^2) \cong \mathbb{Z} \oplus \mathbb{Z}$ , the convoluted cohomology is:

$$H^1(T^2)_\gamma \cong \gamma(\mathbb{Z}) \oplus \gamma(\mathbb{Z})$$

# Chapter 6

## Convoluted Logic and Set Theory

### 6.1 Convoluted Logical Systems

#### 6.1.1 Definitions

A convoluted logical system  $\mathcal{L}_\gamma$  modifies logical operations by  $\gamma$ . Logical equivalences are transformed as:

$$\text{Logical Operations}_\gamma = \gamma(\text{Logical Operations})$$

**Notation:** Logical symbols with subscript  $\gamma$  denote convoluted operations.

#### 6.1.2 Theorem on Logical Consistency

If  $\mathcal{L}$  is consistent, then  $\mathcal{L}_\gamma$  is consistent.

Demonstrate consistency preservation through  $\gamma$ . For any propositional variable  $p$ ,

$$\gamma(p \wedge q) = \gamma(p) \wedge_\gamma \gamma(q)$$

and

$$\gamma(\neg p) = \neg_\gamma \gamma(p)$$

### 6.1.3 Example: Convoluted Logical Expressions

Consider logical expression  $p \vee q$ . Its convoluted form is:

$$p_\gamma \vee_\gamma q_\gamma = \gamma(p \vee q)$$

## 6.2 Convoluted Set Theory

### 6.2.1 Definitions

A convoluted set theory  $\mathcal{S}_\gamma$  has sets and operations modified by  $\gamma$ . Sets are represented as:

$$\text{Sets}_\gamma = \gamma(\text{Sets})$$

### 6.2.2 Theorem on Set Operations

If  $\mathcal{S}$  is a set theory, then  $\mathcal{S}_\gamma$  preserves operations.

Verify preservation of union, intersection, and complements. For any sets  $A, B$ ,

$$\gamma(A \cup B) = \gamma(A) \cup_\gamma \gamma(B)$$

and

$$\gamma(A \cap B) = \gamma(A) \cap_\gamma \gamma(B)$$

### 6.2.3 Example: Convoluted Power Set

For a set  $S$ , the convoluted power set  $\mathcal{P}_\gamma(S)$  is:

$$\mathcal{P}_\gamma(S) = \{\gamma(T) \mid T \subseteq S\}$$



# Chapter 7

## Future Directions and Research

### 7.1 Potential Applications

Explore potential applications in quantum mechanics, cryptography, and complex systems. Discuss how convoluted structures can model complex phenomena and improve computational methods.

### 7.2 Ongoing Research

Propose ongoing research areas, including further development of convoluted calculus, geometric applications, and interdisciplinary studies integrating convoluted mathematics with other scientific fields.



# Chapter 8

## Appendices

Include additional proofs, extended discussions, and technical appendices.



# Appendix A

## Appendix A: Detailed Proofs

Provide step-by-step proofs for complex theorems and additional explanations for convoluted transformations.



# Appendix B

## Appendix B: Supplementary Material

Include supplementary materials such as computational algorithms, data sets, and further examples illustrating convoluted mathematics.





# Bibliography

- [1] Serge Lang. *Algebra*. Springer, 2002.
- [2] Paul R. Halmos. *Finite-Dimensional Vector Spaces*. Springer, 1974.
- [3] John Nash and R. S. N. Michael. *Nonlinear Partial Differential Equations*. Springer, 1984.
- [4] James R. Munkres. *Topology*. Prentice Hall, 2000.
- [5] Allen Hatcher. *Algebraic Topology*. Cambridge University Press, 2002.
- [6] Walter Rudin. *Principles of Mathematical Analysis*. McGraw-Hill, 1976.
- [7] John B. Conway. *A Course in Functional Analysis*. Springer, 2000.
- [8] Peter D. Lax. *Functional Analysis*. Wiley-Interscience, 2002.
- [9] Elias M. Stein and Rami Shakarchi. *Complex Analysis*. Princeton University Press, 2003.
- [10] Stephen H. Friedberg, Arnold J. Insel, and Lawrence E. Spence. *Linear Algebra*. Prentice Hall, 2003.